

27
Feb 08

X - R A Y S O L A R F L A R E S

FEBRUARY 2008

		Start	Max	End			NOAA/ USAF	CMP	Dur	Imp	Total Integrated	Total	Total(3)
Sta	Day	(UT)	(UT)	(UT)	Lat	CMD	Region	Mo	Day (Min)	Xray	Flux(1)	Area(2)	Intensity

NO EVENTS

Note 1: Total integrated flux computed from the event start time to end if available (units=J/m*2).

Note 2: Total area is derived from SXI imagery in units of squared arc seconds of the largest flaring area.

Note 3: Total intensity is derived from SXI imagery in units of data numbers/second of the largest flaring area.

=====

TABLE FORMAT CHANGE: Data are from the GOES full disk xray monitor supplemented with Solar Xray Imager (SXI) from January, 2004, to April 12, 2007. Positions, areas, and intensities are taken from SXI imagery using the largest flare event on the disk. Only the largest event is selected during multiple flares on the disk.

IMPORTANT NOTE: The xray sensor on GOES 12 was turned off on April 12, 2007, at 2250UT. The GOES SXI instrument is also inoperative. GOES 11 is now primary with GOES 10 backup for xray data. Effective April 13, 2007, xray flare locations will be determined by optical flare reports. Xray event times will still be from the xray data.